

Du
30
MARS.
2019

16h00

-
17h00

SÉMINAIRE BOURBAKI

Adam HARPER — The Riemann zeta function in short intervals

Institut Henri Poincaré
Amphithéâtre Hermite
11 rue Pierre-et-Marie-Curie, 75005 Paris

INSCRIPTION

A classical idea for studying the behaviour of complicated functions, like the Riemann zeta function $\zeta(s)$, is to investigate averages of them. For example, the integrals over $T \leq t \leq 2T$ of various powers of $\zeta(1/2+it)$, sometimes multiplied by some other cleverly chosen function, have been investigated extensively to deduce upper and lower bounds for the maximum size of $\zeta(1/2+it)$. More recently, Fyodorov and Keating have proposed the investigation of much shorter integrals over $T \leq t \leq T+1$. This turns out to lead to interesting connections between various issues in number theory, analysis, mathematical physics and probability, such as branching random walk and multiplicative chaos. I will try to explain some of these connections, ideas from the proofs, and what they tell us about the zeta function.



INSTITUT HENRI POINCARÉ - UAR839

Sorbonne Université / CNRS
11 rue Pierre et Marie Curie
75231 Paris Cedex 05

HORAIRES

L'institut :

- lundi au vendredi de 8h30 à 18h,
- fermé les jours fériés.

Le musée - Maison Poincaré :

- lundi, mardi, jeudi et vendredi
de 9h30 à 17h30,
- samedi de 10h à 18h,
- fermé le mercredi et le dimanche.